

A QUICK SURVEY OF FOLIATIONS ON 4-MANIFOLDS

ALEXANDRU SCORPAN

ABSTRACT. We present a few general results on foliations of 4-manifolds by surfaces: existence, tautness, relations to minimal genus of embedded surfaces; as well as some open problems. We hope to stimulate interest in this area.

CONTENTS

1. Introduction	1
2. Generalities	3
3. Thurston's h -principle	5
4. Taut foliations	5
5. More differential geometry	9
6. Minimal genus of embedded surfaces	9
7. Foliations and Gromov compactness	11
8. Singular foliations	14
9. Appendix	18
References	20

1. INTRODUCTION

While there is a wealth of known facts concerning codimension-1 foliations (and in particular foliations on 3-manifolds), the geometric results are quite scarce once one considers higher codimensions, for example 2-dimensional foliations on 4-manifolds. And while there are strong suggestions that the field might yield remarkable insights, one often accuses a lack of methods for attacking it.

This survey has two goals: to quickly gather the known (geometric-topologic) results concerning 2-dimensional foliations on 4-manifolds, and also to frame the context of possible future developments. The biases of the author will become obvious soon enough.

Date: February 7, 2003 (revised February 1, 2008).

2000 Mathematics Subject Classification. Primary 57R30; Secondary 57N13.

Key words and phrases. foliation, four-manifold.

1.1. Quick definition. In general, a k -dimensional *foliation* $\mathcal{F}$ is a partition of a manifold M^m into “slices” called *leaves*. Locally, a foliation must look like the partition of $\mathbb{R}^m$ into $\mathbb{R}^k \times \{0\}$ (see Figure 1).




FIGURE 1. Local model for a foliation

Example 1.1. An important example is the *Reeb foliation* of the solid torus. The boundary torus is a leaf, and the interior is foliated by leaves that are planes. (See Figure 2.)




FIGURE 2. Reeb foliation

For a rigorous introduction to foliations, see for example [CC00]. For a general survey of 4-manifold theory, see [GS99].

Unless otherwise specified, all foliations considered from now on are oriented 2-dimensional foliations of oriented closed 4-manifolds.

1.2. Why bother? One hint that foliations on 4-manifolds are worth studying (especially *taut* foliations) is Kronheimer’s Theorem 6.2 and Conjecture 6.3. Taut foliations might offer *minimal genus bounds* for embedded surfaces.

In a slightly larger context, the relationship (if any) between foliations on 4-manifolds and *Seiberg–Witten theory* is worth elucidating.

Another question worth asking is: For what foliations is the induced almost-complex structure “nice” (*i.e.* close to symplectic). One such problem asks for which foliations does the induced almost-complex structure have *Gromov compactness* (*i.e.* whether the space of J -holomorphic curves of a fixed genus and homology class is compact).

In general, one can hope that foliations will help better visualize, manipulate and understand *almost-complex structures*, maybe in a manner similar to the one in which open-book decompositions help understand contact structures on 3-manifolds.

1.3. Larger context. In Figure 3 we have tried to map the larger mathematical neighborhood in which 2-dimensional foliations on 4-manifolds could be placed:

Foliations, in the presence of a metric, induce *almost-complex structures*. Almost-complex structures are also induced from *symplectic structures*, and are themselves a particular case of a *$\text{spin}^{\mathbb{C}}$ -structure* (one can think of a $\text{spin}^{\mathbb{C}}$ -structure as an almost-complex structure defined on the 2-skeleton). $\text{Spin}^{\mathbb{C}}$ -structures are used to build the *Seiberg–Witten invariants*: one can think of the latter as invariants of various $\text{spin}^{\mathbb{C}}$ -structures. In the case of a symplectic manifold, the Seiberg–Witten invariant can be interpreted as a count of holomorphic curves. The existence of symplectic structures is essentially equivalent to the existence of *Lefschetz pencils* on the manifold. The latter can be thought of as being singular *taut foliations*. In general, almost-complex structures can be deformed to admit singular foliations, see Section 8. Seiberg–Witten theory yields bounds on the *minimal genus* of an embedded surface in a 4-manifold. In certain special cases, taut foliations also offer genus bounds, and it is conjectured that that might be true in general.

2. GENERALITIES

Let $\mathcal{F}$ be a 2-dimensional foliation on a 4-manifold M . It has a tangent bundle $T_{\mathcal{F}}$ (tangent to the leaves), and a normal bundle $N_{\mathcal{F}} =$




FIGURE 3. Context

$T_M/T_{\mathcal{F}}$. An auxiliary Riemannian metric g determines an embedding of the normal bundle $N_{\mathcal{F}}$ as a subbundle of T_M . We then have the g -orthogonal splitting $T_M = T_{\mathcal{F}} \oplus N_{\mathcal{F}}$.

In the presence of a Riemannian metric, a foliation $\mathcal{F}$ induces an *almost-complex structure* $J = J_{\mathcal{F}}$ on M : essentially J rotates the plane-fibers of $T_{\mathcal{F}}$ and $N_{\mathcal{F}}$ by $\pi/2$ in the right directions. Or: pick an orienting orthonormal frame $\{\tau_1, \tau_2, \nu_1, \nu_2\}$ in T_M such that $\tau_1, \tau_2 \in T_{\mathcal{F}}$ and $\nu_1, \nu_2 \in N_{\mathcal{F}}$, and define the almost-complex structure J by $J\tau_1 = \tau_2$ and $J\nu_1 = \nu_2$. The leaves of $\mathcal{F}$ are then J -holomorphic, and the homotopy class of $J_{\mathcal{F}}$ is independent of g .

Due to this almost-complex structure $J_{\mathcal{F}}$, we have well-defined Chern classes: $c_1(T_M, J) = c_1(T_{\mathcal{F}}) + c_1(N_{\mathcal{F}})$ and $c_2(T_M, J) = c_1(T_{\mathcal{F}}) \cdot c_1(N_{\mathcal{F}})$. That can be written:

$$\begin{aligned} c_1(T_M, J) &= e(T_{\mathcal{F}}) + e(N_{\mathcal{F}}) \\ \chi(M) &= e(T_{\mathcal{F}}) \cdot e(N_{\mathcal{F}}) \end{aligned}$$

The class $c_1(T_M)$ also has the *a priori* properties of any Chern class:

$$c_1(T_M) \equiv w_2(M) \pmod{2}$$

$$p_1(M) = c_1(T_M)^2 - 2\chi(M)$$

(Note that $p_1(M) = 3\sigma(M)$, where $\sigma(M)$ is the signature of M .)

3. THURSTON'S h -PRINCIPLE

W. Thurston proved in [Thu74] that any 2-plane field on a 4-manifold can be homotoped to be tangent to a foliation. Namely:

Theorem 3.1 (W. Thurston 1974). *Let τ be a 2-plane field on a manifold M of dimension at least 4. Let K be a compact subset of M such that τ is completely integrable in a neighborhood of K (K can be empty). Then τ is homotopic rel K to a completely integrable plane field.*

This theorem is also true on 3-manifolds (see [Thu76]), but *not* in a relative version.

The theorem is proved by first lifting the plane field to a Haefliger structure, and then deforming the latter to become a foliation using the main theorem of [Thu74]. The latter result has an alternate proof in [EM98].

Of course, as a precondition of the existence of any foliation on M , the tangent bundle of M must split in a sum of two plane bundles (candidates for $T_M = T_{\mathcal{F}} \oplus N_{\mathcal{F}}$). In particular, M must admit almost-complex structures. Nonetheless, if we admit foliations with singularities, these restrictions can be circumvented.

Remark 3.2. By construction, the foliations yielded by Thurston's theorem are not taut: they contain Reeb components. Flexibility comes with a price.

4. TAUT FOLIATIONS

(Most statements of this section admit obvious generalizations to arbitrary dimensions and codimensions.)

4.1. Definition. A foliation $\mathcal{F}$ on a Riemannian manifold (M, g) is called *minimal* if all its leaves are minimal surfaces in (M, g) (*i.e.* they locally minimize area; for any compact piece K of a leaf, any small perturbation of K rel ∂K will have bigger area; that is equivalent to each leaf having zero mean curvature).

A foliation $\mathcal{F}$ on M is called *taut* if there is a Riemannian metric g such that $\mathcal{F}$ is minimal in (M, g)

Remark 4.1. In the special case of a codimension-1 foliation, tautness is equivalent to the existence of a 1-manifold transverse to $\mathcal{F}$ and crossing all the leaves. A similar condition is much too strong for higher codimensions.

4.2. Mean curvature and characteristic form. We define the *mean curvature* vector field $H_{\mathcal{F},g}$ as (half) the normal component of $\nabla_{\tau_1}^g \tau_1 + \nabla_{\tau_2}^g \tau_2$, for any orthonormal orienting frame $\{\tau_1, \tau_2\}$ in $T_{\mathcal{F}}$, with ∇^g the Levi-Civita connection of g . Perturbing a leaf in the direction opposite to H most decreases its area. If $H = 0$ along a leaf, then that leaf is a minimal surface in (M, g) .

Proposition 4.2. *Let $\mathcal{F}$ be a foliation on (M, g) . Let μ be any 2-form on M such that $\mu|_{\text{Leaf}} = \text{vol}_g|_{\text{Leaf}}$ and $\mu(\tau, \nu) = 0$ for any $\tau \in T_{\mathcal{F}}$ and $\nu \in N_{\mathcal{F}}$. Denote by H the mean curvature vector field of the leaves. Then*

$$d\mu(\tau_1, \tau_2, \nu) = \langle H, \nu \rangle$$

for any orthonormal orienting basis τ_1, τ_2 in $T_{\mathcal{F}}$ and any $\nu \in N_{\mathcal{F}}$.

In particular, $\mathcal{F}$ is minimal if and only if $d\mu(\tau_1, \tau_2, \nu) = 0$ for all $\tau_1, \tau_2 \in T_{\mathcal{F}}$ and $\nu \in N_{\mathcal{F}}$. $\square$

Forms μ as above always exist, and are not unique. A particular such μ is the *characteristic form* $\chi = \chi_{\mathcal{F},g}$, defined to be the unique 2-form such that $\chi|_{\text{Leaf}} = \text{vol}_g|_{\text{Leaf}}$ and $\chi(x, \nu) = 0$ for any $x \in T_M$ and $\nu \in N_{\mathcal{F}}$. (In other words, χ is just the volume form along the leaves.)

The condition $d\mu(\tau_1, \tau_2, \nu) = 0$ is strictly weaker than asking that μ be closed. For example, for the characteristic form:

Lemma 4.3. *If $d\chi_{\mathcal{F},g} = 0$, then $\mathcal{F}$ is minimal and $\mathcal{F}^\perp$ is integrable.* $\square$

The integrability of $\mathcal{F}^\perp$ is measured by the other significant component of $d\chi$, namely $d\chi(\tau, \nu_1, \nu_2)$ for $\tau \in T_{\mathcal{F}}$ and $\nu_1, \nu_2 \in N_{\mathcal{F}}$. More precisely, we have

$$d\chi(\tau, \nu_1, \nu_2) = \chi(\tau, [\nu_1, \nu_2])$$

4.3. Rummmler's criterion. In the direction opposite to the above Proposition 4.2, it was proved in [Rum79]:

Theorem 4.4 (H. Rummmler 1979). *Let $\mathcal{F}$ be a foliation on M . Assume that there is a 2-form μ such that $\mu|_{\text{Leaf}} > 0$ and $d\mu|_{\text{Leaf}} = 0$. Then $\mathcal{F}$ is taut.*

When writing “ $\mu|_{\text{Leaf}} > 0$ and $d\mu|_{\text{Leaf}} = 0$ ” we mean “ $\mu(\tau_1, \tau_2) > 0$ and $d\mu(\tau_1, \tau_2, z) = 0$ for all orienting pairs $\tau_1, \tau_2 \in T_{\mathcal{F}}$ and $z \in T_M$ ”.

Proof. If μ is positive on the leaves, then there is a metric g^τ , defined only on $T_{\mathcal{F}}$ such that $\mu|_{\text{Leaf}} = \text{vol}_{g^\tau}|_{\text{Leaf}}$. We need to extend g^τ to a genuine metric g on M that will determine a normal bundle $N_{\mathcal{F}}$ for $\mathcal{F}$ such that $\mu(\tau, \nu) = 0$ for any $\tau \in T_{\mathcal{F}}$ and $\nu \in N_{\mathcal{F}}$. We will first find $N_{\mathcal{F}}$, then extend g^τ . Concretely, if τ_1, τ_2 is any basis in $T_{\mathcal{F}}$, we define $N_{\mathcal{F}}$ as $\ker \mu(\tau_1, \cdot) \cap \ker \mu(\tau_2, \cdot)$. Then, after choosing some random metric g^ν on $N_{\mathcal{F}}$, we can extend g^τ to the whole $T_M = T_{\mathcal{F}} \oplus N_{\mathcal{F}}$ simply as $g = g^\tau \oplus g^\nu$. In view of Proposition 4.2, this g will make $\mathcal{F}$ minimal. $\square$

Remark 4.5. A condition stronger than tautness would be the existence of a 2-form μ such that $\mu|_{\text{Leaf}} > 0$ and $d\mu = 0$. The foliation $\mathcal{F}$ is then *calibrated* by μ (see [HL82]), and, for a suitable metric g , the leaves are not only locally area minimizing, but are locally area minimizing in their homology classes (*i.e.* if K is any compact piece of a leaf and K' is any immersed surface with $\partial K' = \partial K$ and representing the same class as K in $H_2(M, \partial K)$, then K' has bigger area than K).

Example 4.6. A Reeb component is not taut. Denote by C a solid torus foliated so that the boundary torus ∂C is a leaf, and the interior is foliated by planes (see Figure 2). Consider any 2-dimensional foliation $\mathcal{F}$ (of any codimension) containing such a C . Assume that $\mathcal{F}$ is taut. Then there is a 2-form μ satisfying the conditions from Rummier's Theorem 4.4. But then

$$0 = \int_C d\mu = \int_{\partial C} \mu > 0$$

which is impossible.

The strong link between 2-forms and minimality of foliations is also suggested (via almost-complex structures) by the following formula:

Lemma 4.7. *Let $g(x, y) = \langle x, y \rangle$ be a Riemannian metric on M^4 and ∇ its Levi-Civita connection. Let J be any g -orthogonal almost-complex structure, and let $\omega(x, y) = \langle Jx, y \rangle$ be its fundamental 2-form. Let x, z be any vector fields on M . Then:*

$$d\omega(x, Jx, z) = \langle [x, Jx], Jz \rangle - \langle \nabla_x x + \nabla_{Jx} Jx, z \rangle$$

The term $[x, Jx]$ measures the integrability of the J -holomorphic plane field $\mathbb{R}\langle x, Jx \rangle$, while the normal component of the term $\nabla_x x + \nabla_{Jx} Jx$ is the mean curvature of the plane field $\mathbb{R}\langle x, Jx \rangle$, and thus measures its g -minimality. (Lemma 4.7 will be proved at the end of the paper, in Section 9.)

4.4. Sullivan’s criterion. In what follows, we wish to at least suggest the nature of Sullivan’s criterion for tautness. Rigor must suffer. In a way, Sullivan’s criterion is a far-reaching generalization of the Reeb Example 4.6 above.

A 2-chain σ in M is called a *tangent homology* in $(M, \mathcal{F})$ if there is a 3-chain ξ such that $\partial\xi = \sigma$ and $T_{\mathcal{F}}|_{\xi} \subset T_{\xi}$. (In other words, σ is a boundary that can be “filled-in” using a chain ξ that is tangent to $\mathcal{F}$.)

Lemma 4.8. *If a sequence of tangent homologies converges to an (oriented) union of compact leaves of $\mathcal{F}$, then $\mathcal{F}$ is not taut.*

Proof. Let $L_1, \dots, L_k$ be some compact leaves of $\mathcal{F}$, and let $\sigma_n = \partial\xi_n$ be a sequence of tangent homologies that converges to $L_1 \cup \dots \cup L_k$. Assume that $\mathcal{F}$ is taut, and thus there is a 2-form μ with $\mu|_{\text{Leaf}} > 0$ and $d\mu|_{\text{Leaf}} = 0$. Then:

$$0 = \int_{\xi_n} d\mu = \int_{\partial\xi_n} \mu = \int_{\sigma_n} \mu \xrightarrow{n} \int_{L_1 \cup \dots \cup L_k} \mu > 0$$

which is impossible □

(The appropriate notion of convergence here is, of course, a convergence “in measure”.)

Of course, in general, $\mathcal{F}$ might not have compact leaves at all. Nonetheless, a set of non-compact leaves might come back on itself in a regular enough fashion that it could be called a “foliation cycle”. Somewhat more rigorously, a *foliation cycle* can be defined as a transverse measure that is holonomy-invariant. Think of this measure as a way of weighting the leaves of $\mathcal{F}$. Then:

Theorem 4.9 (D. Sullivan 1979). *A foliation $\mathcal{F}$ is taut if and only if no non-trivial foliation cycle can be approximated by tangent homologies.* □

This result was proved in [Sul79], based on the theory developed in [Sul76]. A complete recent discussion can be found in [CC00, Ch. 10]. The rigorous development of these ideas involves currents.

A (rarely applicable) consequence of this criterion is:

Corollary 4.10. *Let $\mathcal{F}$ be a foliation on M . If for every leaf L of $\mathcal{F}$ there is a surface transverse to $\mathcal{F}$ that crosses L , then $\mathcal{F}$ is taut. In fact, $\mathcal{F}$ can be calibrated.*

Surfaces transverse to foliations are a rare occurrence, though (unlike the codimension-1 case, where tautness is equivalent to the existence of transverse circles).

5. MORE DIFFERENTIAL GEOMETRY

Here are a couple of general results of a differential geometric nature, from [Bri84] and [GN95]:

Proposition 5.1 (B. Fabiano 1984). *Let $\mathcal{F}$ be a minimal codimension-2 foliation on the Riemannian manifold M^m . If M has non-negative Ricci curvature and $\mathcal{F}$ admits a complementary foliation $\mathcal{F}^\perp$, then: $\mathcal{F}$ is totally geodesic, or $e(\mathcal{F}^\perp) \neq 0$*

Proposition 5.2 (F.J. García & A.M. Naveira 1995). *Let $\mathcal{F}$ be a minimal foliation on the Riemannian manifold M^m . If M has non-positive Ricci curvature and $\mathcal{F}$ admits a complementary foliation $\mathcal{F}^\perp$, then: $\mathcal{F}$ is totally geodesic, or the leaves of $\mathcal{F}^\perp$ have non-zero scalar curvature.*

As one can notice, these general results have pretty restrictive conditions (mainly: the existence of a complementary foliation). In general, it is worth noticing that foliations of codimension ≥ 2 are much harder to study than those of codimension 1. This, in part, explains the success of foliation theory for investigating 3-manifolds, and the lack of much of a theory for 4-manifolds.

6. MINIMAL GENUS OF EMBEDDED SURFACES

Given any class $a \in H_2(M; \mathbb{Z})$, there are always embedded surfaces in M that represent it. An open problem is to determine how simple such surfaces can be, or, in other words, what is the minimal genus that a surface representing a can have. (Remember that $\chi(S) = 2 - 2g(S)$, so minimum genus is maximum Euler characteristic.)

6.1. Adjunctions. Notice that, if S is a J -holomorphic surface for some almost-complex structure J , then

$$\chi(S) + S \cdot S = c_1(J) \cdot S$$

(simply because $c_1(J) \cdot S = c_1(T_M|_S) = c_1(T_S) + c_1(N_S) = \chi(S) + S \cdot S$). This equality is known as the *Adjunction formula* for S .

In general, the main and most powerful tool for obtaining genus bounds comes from Seiberg–Witten theory [KM94, OS00]:

Proposition 6.1 (Adjunction Inequality). *Let S be any embedded surface in M . Assume that either M is of Seiberg–Witten simple type and S has no sphere components, or that $S \cdot S \geq 0$. Then, for any Seiberg–Witten basic class ϵ , we have:*

$$\chi(S) + S \cdot S \leq \epsilon \cdot S$$

In particular, if J is an almost-complex structure admitting a symplectic structure, then

$$\chi(S) + S \cdot S \leq c_1(J) \cdot S$$

Notice that one can reverse the orientation of S and change only the sign of $\epsilon \cdot S$. Thus one often writes the adjunction inequality more sharply as

$$\chi(S) + S \cdot S \leq -|\epsilon \cdot S|$$

6.2. Enter foliation. Nonetheless, the bounds offered by Seiberg–Witten basic classes are not always sharp. For example, in the case of manifolds $M = N^3 \times \mathbb{S}^1$, P. Kronheimer has proved in [Kro99] that foliations give *better* bounds:

Theorem 6.2 (P. Kronheimer 1999). *Consider $M = N^3 \times \mathbb{S}^1$, with N a closed irreducible 3-manifold. Let $\overline{\mathcal{F}}$ be a taut foliation in N , with Euler class $\bar{\epsilon} = e(T_{\overline{\mathcal{F}}})$. Let ϵ be the image of $\bar{\epsilon}$ in $H^2(N \times \mathbb{S}^1)$. Then, for any embedded surface S in M , without sphere components, we have*

$$\chi(S) + S \cdot S \leq \epsilon \cdot S$$

In general, ϵ is *not* a Seiberg–Witten basic class. (Nonetheless, the proof of Theorem 6.2 does use Seiberg–Witten theory: the taut foliation $\overline{\mathcal{F}}$ is perturbed to a tight contact structure, which is then symplectically filled in a suitable way, and a version of the Seiberg–Witten invariants is used.)

A taut foliation $\overline{\mathcal{F}}$ on N^3 induces an obvious product foliation $\mathcal{F} = \overline{\mathcal{F}} \times \mathbb{S}^1$ on $M = N \times \mathbb{S}^1$ (with leaves $\text{Leaf} \times \{pt\}$). Then $\mathcal{F}$ is also taut (pick a product metric), $\epsilon = e(T_{\mathcal{F}})$ is the pull-back of $\bar{\epsilon} = e(T_{\overline{\mathcal{F}}})$, and the almost-complex structure that $\mathcal{F}$ determines has $c_1(J_{\mathcal{F}}) = \epsilon = e(T_{\mathcal{F}})$. One can then try to generalize Theorem 6.2 as

Conjecture 6.3 (P. Kronheimer 1999). *Let $\mathcal{F}$ be a taut foliation on M^4 , and $J_{\mathcal{F}}$ be an almost-complex structure induced by $\mathcal{F}$. Then, for any embedded surface S without sphere components, we have*

$$\chi(S) + S \cdot S \leq c_1(J_{\mathcal{F}}) \cdot S$$

A few extra requirements are needed, *e.g.* to exclude manifolds like $S \times \mathbb{S}^2$. Kronheimer also proposes that the foliations be allowed singularities.

Remark 6.4. The situation in Theorem 6.2 has another peculiarity: $\mathcal{F}$ admits transverse foliations. Indeed, since $\overline{\mathcal{F}}$ has codimension 1, any nowhere-zero vector-field in N^3 normal to $\overline{\mathcal{F}}$ integrates to a 1-dimensional foliation of N that is transverse to $\overline{\mathcal{F}}$. By multiplying

its leaves by $\mathbb{S}^1$, this 1-dimensional foliation induces a 2-dimensional foliation in M that is transverse to $\mathcal{F}$.

One could thus think of strengthening the hypothesis of Conjecture 6.3 by requiring $\mathcal{F}$ not only to be taut, but also to admit a transverse foliation. One might push things even further and ask that the second foliation be taut as well. But then one almost runs into:

Proposition 6.5 (Two taut makes one symplectic). *Let $\mathcal{F}$ and $\mathcal{G}$ be transverse foliations on M^4 . If there is a metric g that makes both $\mathcal{F}$ and $\mathcal{G}$ be minimal and orthogonal, then M must admit a symplectic structure. Therefore, for any embedded surface S we have*

$$\chi(S) + S \cdot S \leq c_1(J_{\mathcal{F}}) \cdot S$$

(This is an immediate consequence of Lemma 4.7.)

Remark 6.6. No taut, no symplectic. At the other extreme, if M admits a *non-taut* foliation $\mathcal{F}$, then *no* almost-complex structure compatible with $\mathcal{F}$ admits symplectic structures (or, more directly, M admits no symplectic structures making the leaves of $\mathcal{F}$ symplectic submanifolds).

6.3. Contrast with dimension 3. Taut foliations on 3-manifolds are well-understood, and are strongly related to minimal genus surfaces [Thu86, Gab83]:

Proposition 6.7 (W. Thurston 1986). *Let N^3 be a closed irreducible 3-manifold, and $\mathcal{F}$ a taut foliation on N . For any embedded surface S without sphere-components, we have*

$$\chi(S) \leq e(T_{\mathcal{F}}) \cdot S$$

(Usually written more sharply as $\chi(S) \leq -|e(T_{\mathcal{F}}) \cdot S|$.)

Proposition 6.8 (D. Gabai 1983). *Let N^3 be a closed irreducible 3-manifold. An embedded surface S has minimal genus if and only if it is the leaf of a taut foliation of N .* $\square$

No similar statement is known for 4-manifolds. For a survey of foliations on 3-manifolds, see for example [Gab01].

7. FOLIATIONS AND GROMOV COMPACTNESS

First, an example that shows the flexibility of foliations:

Example 7.1. Creating a torus leaf. Let $\mathcal{F}$ be *any* foliation on a 4-manifold M . Let $c : \mathbb{S}^1 \rightarrow M$ be any embedding. The curve c can always be slightly perturbed to be transverse to $\mathcal{F}$. Choose another local coordinate near c , transverse both to $\mathcal{F}$ and to c , and think of




FIGURE 4. Creating a torus leaf (3D movie)

it as *time* (with c appearing at time $t = 0$). Start at time $t = -1$. As time goes on, begin pushing more and more the leaves of $\mathcal{F}$ parallel with the direction of c , wrapping them around more and more as time approaches $t = 0$ (see Figures 4 and 5). At $t = 0$, we can fit in a torus leaf, with the interior of the torus foliated by leaves diffeomorphic to $\mathbb{R}^2$ —a *Reeb component*. As time goes on from $t = 0$, we play the movie backward. Notice that the new foliation is *homotopic* with the one we started with (compare with Corollary 8.8).

Combining with Theorem 8.2 below, this yields:

Corollary 7.2. *Any almost-complex structure is homotopic to one for which Gromov compactness fails.*




FIGURE 5. Creating a torus leaf (2D movie)

Gromov compactness here means the compactness of the space of all J -holomorphic curves (curve = real surface). In other words, any sequence of holomorphic curves $f_n : (\Sigma, j_n) \rightarrow (M, J)$ has a subsequence converging to a limit $f : \Sigma^* \rightarrow M$ that is J -holomorphic (and may have nodal singularities, see also Figure 6; the limit domain Σ^* is obtained by collapsing circles of Σ). Gromov compactness always holds for symplectic structures, due to the following [Gro85]:

Theorem 7.3 (Gromov compactness). *Let J be an almost-complex structure on M , and g a compatible Riemannian metric. Let $f_n : (\Sigma, j_n) \rightarrow (M, J)$ be a sequence of holomorphic compact curves in M . If the g -area of $f_n(\Sigma)$ is bounded, then f_n has a subsequence that converges to a holomorphic curve $f : (\Sigma^*, j) \rightarrow (M, J)$, which might have nodal singularities.*

And the area bound condition is satisfied if J admits a compatible symplectic structure, and if all $f_n(\Sigma)$ represent a same homology class. (This is actually how the theorem is most used.) For a thorough discussion, see for example [AL94].




FIGURE 6. Apparition of singularities in the limit

Proof of Corollary 7.2. By Theorem 8.2, an almost-complex structure can be deformed till there is a singular foliation $\mathcal{F}$ with all leaves holomorphic. As in Example 7.1 above, create a torus leaf. Actually, by “freezing” the movie at $t = 0$ (expanding the frame at $t = 0$ to all $t \in [-\varepsilon, \varepsilon]$), create a lot of tori. Now pick a second curve, orthogonal to these tori, and apply that example again. What appears in the end is a torus that explodes to make room for a new Reeb component. Thinking in terms of an almost-complex structure J induced by the final foliation, we have a sequence of J -holomorphic tori that has no decent limit. $\square$

Question 7.4 (R. Kirby 2002). *Let $\mathcal{F}$ be a foliation on M , and J an almost-complex structure making the leaves J -holomorphic. What conditions imposed on $\mathcal{F}$ insure that Gromov compactness holds for J ?*

In the extreme, if $\mathcal{F}$ is a Lefschetz pencil (see Example 8.1), then Gromov compactness holds (the manifold is symplectic). Compare also with Proposition 6.5.

The flexibility from our examples, at least, is done away with if we require the foliations to be *taut* (since that excludes Reeb components).

8. SINGULAR FOLIATIONS

For a fuller treatment of the topic of this section, see [Sco03].

For a foliation $\mathcal{F}$ to exist on a manifold M , the tangent bundle must split $T_M = T_{\mathcal{F}} \oplus N_{\mathcal{F}}$. Since in general that does not happen, one must allow for singularities of $\mathcal{F}$. An important example is [Don99]:

Theorem 8.1 (S.K. Donaldson). *Let J be an almost-complex structure on M that admits a compatible symplectic structure. Then J can*




FIGURE 7. A Lefschetz pencil

be deformed to an almost-complex structure J' such that M admits a Lefschetz pencil with J' -holomorphic fibers.

A **Lefschetz pencil** is a *singular fibration* $M \rightarrow \mathbb{CP}^1$ (see Figures 7 and 8) with singularities modeled locally by

$$(z_1, z_2) \mapsto z_1/z_2 \quad \text{or} \quad (z_1, z_2) \mapsto z_1 z_2$$

for suitable local complex coordinates (compatible with the orientation of M). Note that all fibers pass through all singularities of type z_1/z_2 . The existence of a Lefschetz pencil is equivalent to the existence of a symplectic structure. See [GS99, ch. 8] for a survey.

One can think of a Lefschetz pencil as a special case of a taut *singular foliation*.

As mentioned before, a (non-singular) foliation $\mathcal{F}$ on M induces almost-complex structures, and we have $c_1(J_{\mathcal{F}}) = e(T_{\mathcal{F}}) + e(N_{\mathcal{F}})$ and $\chi(M) = e(T_{\mathcal{F}}) \cdot e(N_{\mathcal{F}})$. If the foliation $\mathcal{F}$ has singularities, then the




FIGURE 8. Singularities in a Lefschetz pencil

second equality above fails, and the defect $\chi(M) - e(T_{\mathcal{F}}) \cdot e(N_{\mathcal{F}})$ measures the number of singularities (or, for more general singularities, their complexity).

Call a class $c \in H^2(M; \mathbb{Z})$ a *complex class* of the 4-manifold M if

$$c \equiv w_2(M) \pmod{2} \quad \text{and} \quad p_1(M) = c^2 - 2\chi(M)$$

An element $c \in H^2(M; \mathbb{Z})$ is a complex class if and only if there is an almost-complex structure J on M such that $c_1(T_M, J) = c$.

Theorem 8.2 (Existence theorem). *Let $c \in H^2(M; \mathbb{Z})$ be a complex class, and let $c = \tau + \nu$ be any splitting such that $\chi(M) - \tau\nu \geq 0$. Then there is a singular foliation $\mathcal{F}$ with $e(T_{\mathcal{F}}) = \tau$ and $e(N_{\mathcal{F}}) = \nu$, with $n = \chi(M) - \tau\nu$ singularities that can be chosen to be modeled on the levels of the complex functions $(z_1, z_2) \mapsto z_1/z_2$ or $(z_1, z_2) \mapsto z_1 z_2$.*

Remark 8.3. Due to the singularities, the bundles $T_{\mathcal{F}}$ and $N_{\mathcal{F}}$ are only defined on $M \setminus \{\text{singularities}\}$. Their Euler classes *a priori* belong to $H^2(M \setminus \{\text{singularities}\}; \mathbb{Z})$, but can be pulled-back to $H^2(M; \mathbb{Z})$, since the isolated singularities can be chosen to affect only the 4-skeleton of M , and thus to not influence H^2 .

Remark 8.4. Unlike a Lefschetz pencil, in general *not all* leaves of the foliation pass through the z_1/z_2 -singularities. If they did, then the foliation would be taut, which is rather rare.

Finding a splitting $c = \tau + \nu$ with $\chi(M) - \tau\nu \geq 0$ is possible for most 4-manifolds that admit almost-complex structures. For example,

if

$$\chi(M) \geq 0$$

(e.g. for all *simply-connected* M 's), then one can choose either one of τ or ν to be 0, and conclude that such foliations exist. Or:

Lemma 8.5. *If $b_2^+(M) > 0$, then there are infinitely many splittings $c = \tau + \nu$ with $\chi(M) - \tau\nu \geq 0$ (and thus infinitely many homotopy types of foliations). If $b_2^+(M) = 0$, there are at most finitely many.*

Proof. If $b_2^+(M) > 0$, there is a class α with $\alpha \cdot \alpha > 0$. Choose $\tau = c - k\alpha$ and $\nu = k\alpha$ ($k \in \mathbb{Z}$). Then $\chi(M) - \tau\nu = \chi(M) - kc\alpha + k^2\alpha^2$, and for k big enough it will be positive. $\square$

The main restriction to the existence of such foliations remains, of course, the existence of a complex class. But Theorem 8.2 can be generalized for the case when c is merely an *integral lift* of $w_2(M)$ (corresponding to weakening almost-complex structures to *spin^C-structures*). In that case, singularities are also modeled using local complex coordinates, but are allowed to be compatible either with the orientation of M or with the *opposite orientation*. This is similar to the generalization of Lefschetz pencils to *achiral Lefschetz pencils*, see [GS99, §8.4].

The singularities of $\mathcal{F}$ are exactly the singularities that appear in a Lefschetz pencil. They can be chosen in either combination of types as long as their number is $n = \chi(M) - \tau\nu$. For example, there are always foliations with only z_1/z_2 -singularities, that can thus be eliminated by blowing-up. In fact, other choices of singularities are possible.

Namely, for any isolated singularity p of a foliation that is compatible with an almost-complex structure one can define its *Hopf degree* $\deg p \geq 0$ (essentially a Hopf invariant of the tangent plane field above a small 3-sphere around p), and then one can choose any suitable set of singularities $\{p_i\}$ as long as

$$\sum \deg p_i = \chi(M) - \tau\nu$$

Let $\mathcal{F}$ be a foliation. If S is a closed transversal of $\mathcal{F}$, then we must have

$$\begin{aligned} e(T_{\mathcal{F}}) \cdot S &= e(T_{\mathcal{F}}|_S) = e(N_S) = S \cdot S \\ e(N_{\mathcal{F}}) \cdot S &= e(N_{\mathcal{F}}|_S) = e(T_S) = \chi(S) \end{aligned}$$

These conditions are, in fact, sufficient:

Theorem 8.6 (Closed transversal). *Let S be a closed connected surface. Let c be a complex class with a splitting $c = \tau + \nu$ such that $\chi(M) - \tau\nu \geq 0$. If*

$$\chi(S) = \nu \cdot S \quad S \cdot S = \tau \cdot S$$

then there is a singular foliation $\mathcal{F}$ with $e(T_{\mathcal{F}}) = \tau$, $e(N_{\mathcal{F}}) = \nu$, and having S as a closed transversal.

If, on the other hand, S is a closed leaf of $\mathcal{F}$, then we have

$$\begin{aligned} e(T_{\mathcal{F}}) \cdot S &= e(T_{\mathcal{F}}|_S) = e(T_S) = \chi(S) \\ e(N_{\mathcal{F}}) \cdot S &= e(N_{\mathcal{F}}|_S) = e(N_S) = S \cdot S \end{aligned}$$

Conversely:

Theorem 8.7 (Closed leaf). *Let S be a closed connected surface with $S \cdot S \geq 0$. Let c be a complex class with a splitting $c = \tau + \nu$ such that $\chi(M) - \tau\nu \geq S \cdot S$. If*

$$\chi(S) = \tau \cdot S \quad S \cdot S = \nu \cdot S$$

then there is a singular foliation $\mathcal{F}$ with $e(T_{\mathcal{F}}) = \tau$, $e(N_{\mathcal{F}}) = \nu$, and having S as a closed leaf. (The number of singularities along S is $S \cdot S$.)

An immediate consequence is:

Corollary 8.8 (Trivial tori). *A homologically-trivial torus can always be made a leaf or a transversal of a foliation.*

Such flexibility is a strong suggestion that more rigidity is needed in order to actually catch any of the topology of M with the aid of foliations. Requiring foliations to be taut seems, once again, a natural suggestion.

9. APPENDIX

9.1. Minimality and 2-forms.

Proof of Lemma 4.7. We prove that, if g is a Riemannian metric, ∇ its Levi-Civita connection, J be any g -orthogonal almost-complex structure, and $\omega(x, y) = \langle Jx, y \rangle$ its fundamental 2-form, then, for any vector fields x, z on M , we have:

$$(d\omega)(x, Jx, z) = \langle [x, Jx], Jz \rangle - \langle \nabla_x x + \nabla_{Jx} Jx, z \rangle$$

For any 2-form α we have:

$$\begin{aligned} (d\alpha)(x, y, z) &= (\nabla_x \alpha)(y, z) + (\nabla_y \alpha)(z, x) + (\nabla_z \alpha)(x, y) \\ &= x\alpha(y, z) + y\alpha(z, x) + z\alpha(x, y) \\ &\quad - \alpha(\nabla_x y, z) - \alpha(\nabla_y z, x) - \alpha(\nabla_z x, y) \\ &\quad - \alpha(y, \nabla_x z) - \alpha(z, \nabla_y x) - \alpha(x, \nabla_z y) \end{aligned}$$

Applying this to $\omega(a, b) = \langle Ja, b \rangle$, we have:

$$\begin{aligned} (d\omega)(x, Jx, z) &= -x\langle x, z \rangle + (Jx)\langle Jz, x \rangle + z\langle Jx, Jx \rangle \\ &\quad - \langle J\nabla_x Jx, z \rangle - \langle J\nabla_{Jx} z, x \rangle - \langle J\nabla_z x, Jx \rangle \\ &\quad + \langle x, \nabla_x z \rangle - \langle Jz, \nabla_{Jx} x \rangle - \langle Jx, \nabla_z Jx \rangle \end{aligned}$$

Using that $\langle a, Jb \rangle = -\langle Ja, b \rangle$ we get:

$$\begin{aligned} (d\omega)(x, Jx, z) &= -x\langle x, z \rangle + (Jx)\langle Jz, x \rangle + z\langle x, x \rangle \\ &\quad + \langle \nabla_x Jx, Jz \rangle + \langle \nabla_{Jx} z, Jx \rangle - \langle \nabla_z x, x \rangle \\ &\quad + \langle x, \nabla_x z \rangle - \langle Jz, \nabla_{Jx} x \rangle - \langle Jx, \nabla_z Jx \rangle \end{aligned}$$

Since $\langle \nabla_z x, x \rangle = \frac{1}{2}z\langle x, x \rangle$ and $\langle Jx, \nabla_z Jx \rangle = \frac{1}{2}z\langle Jx, Jx \rangle = \frac{1}{2}z\langle x, x \rangle$, we cancel the last terms of each line, and get:

$$\begin{aligned} (d\omega)(x, Jx, z) &= -x\langle x, z \rangle + (Jx)\langle Jz, x \rangle \\ &\quad + \langle \nabla_x Jx, Jz \rangle + \langle \nabla_{Jx} z, Jx \rangle \\ &\quad + \langle x, \nabla_x z \rangle - \langle Jz, \nabla_{Jx} x \rangle \end{aligned}$$

But $x\langle x, z \rangle = \langle \nabla_x x, z \rangle + \langle x, \nabla_x z \rangle$, so $(Jx)\langle Jz, x \rangle = -(Jx)\langle z, Jx \rangle = -\langle \nabla_{Jx} z, Jx \rangle - \langle z, \nabla_{Jx} Jx \rangle$, and therefore:

$$\begin{aligned} (d\omega)(x, Jx, z) &= -\langle \nabla_x x, z \rangle - \langle x, \nabla_x z \rangle - \langle \nabla_{Jx} z, Jx \rangle - \langle z, \nabla_{Jx} Jx \rangle \\ &\quad + \langle \nabla_x Jx, Jz \rangle + \langle \nabla_{Jx} z, Jx \rangle + \langle x, \nabla_x z \rangle - \langle Jz, \nabla_{Jx} x \rangle \\ (d\omega)(x, Jx, z) &= -\langle \nabla_x x, z \rangle - \langle z, \nabla_{Jx} Jx \rangle + \langle \nabla_x Jx, Jz \rangle - \langle Jz, \nabla_{Jx} x \rangle \end{aligned}$$

Since ∇ is torsion-free, we have $\nabla_x Jx - \nabla_{Jx} x = [x, Jx]$, so:

$$(d\omega)(x, Jx, z) = \langle [x, Jx], Jz \rangle - \langle \nabla_x x + \nabla_{Jx} Jx, z \rangle$$

which concludes the proof. $\square$

In particular, if ω is symplectic (*i.e.* $d\omega = 0$), then any integrable J -holomorphic plane field is g -minimal, and, vice-versa, any g -minimal J -holomorphic plane field must be integrable. The converse is also true: If there are enough J -holomorphic integrable minimal plane fields, then ω must be symplectic. Thus:

Corollary 9.1. *Assume that M admits two transversal 2-dimensional foliations $\mathcal{F}$ and $\mathcal{G}$ such that: there is a metric g such that both $\mathcal{F}$ and $\mathcal{G}$ are g -minimal; and there is an g -orthogonal almost-complex structure J that makes both $\mathcal{F}$ and $\mathcal{G}$ be J -holomorphic. Then M admits the symplectic structure $\omega(x, y) = g(Jx, y)$.*

In particular, if the first condition is satisfied, and further $\mathcal{F}$ and $\mathcal{G}$ are g -orthogonal, then the second condition is automatically satisfied, and Proposition 6.5 follows:

If a Riemannian manifold M admits two g -orthogonal and g -minimal foliations, then M admits a symplectic structure.

9.2. Foliated surgery. We add here a few comments on some surgical techniques that can be used to combine foliations.

9.2. Transverse surfaces to foliations can be used to glue foliated manifolds using a *normal connected sum*: Let $(M, \mathcal{F})$ and $(N, \mathcal{G})$ be foliated manifolds. Assume that there is an embedded surface S_1 in M transverse to $\mathcal{F}$, and an embedded surface S_2 in N transverse to $\mathcal{G}$. Assume further that $S_1 \approx S_2$ and $S_1 \cdot S_1 = -S_2 \cdot S_2$. Then the foliated manifolds $(M, \mathcal{F})$ and $(N, \mathcal{G})$ admit a foliated normal connected sum along $S_1 \approx S_2$.

9.3. For regular *connected sums*, one can use pencil singularities of opposite orientations. Namely, when doing the connected sum, pick small 3-spheres around two pencil singularities of opposite types. The result of the connected sum is a foliated manifold with one less positive pencil singularity and one less negative pencil singularity.

9.4. Another surgery is the *logarithmic transformation* performed along a torus T that is transverse to a foliation and has $T \cdot T = 0$. One takes a neighborhood $T \times D^2$ of the torus T , cuts it out and glues it back in using an automorphism of the boundary $\partial(T \times D^2) = \mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1$ (see Figure 9, where a logarithmic transformation of multiplicity 2 is suggested, with the unrepresented $\mathbb{S}^1$ -factor belonging to T). If T is transverse to the foliation $\mathcal{F}$, then one way of continuing the foliation across the glued-in $T \times D^2$ is suggested in Figure 10, but it needs a (circle of) Reeb components and thus ruins tautness.

REFERENCES

- [AL94] Michèle Audin and Jacques Lafontaine (eds.), *Holomorphic curves in symplectic geometry*, Progress in Mathematics, vol. 117, Birkhäuser Verlag, Basel, 1994. MR **95i**:58005




FIGURE 9. Logarithmic transformation (multiplicity 2)

- [Bri84] Fabiano Brito, *A remark on minimal foliations of codimension two*, Tohoku Math. J. (2) **36** (1984), no. 3, 341–350. MR **85m**:53029
- [CC00] Alberto Candel and Lawrence Conlon, *Foliations. I*, Graduate Studies in Mathematics, vol. 23, American Mathematical Society, Providence, RI, 2000. MR **2002f**:57058
- [Don99] S. K. Donaldson, *Lefschetz pencils on symplectic manifolds*, J. Differential Geom. **53** (1999), no. 2, 205–236. MR **2002g**:53154
- [EM98] Y. Eliashberg and N. M. Mishachev, *Wrinkling of smooth mappings. III. Foliations of codimension greater than one*, Topol. Methods Nonlinear Anal. **11** (1998), no. 2, 321–350. MR **2000a**:57073
- [Gab83] David Gabai, *Foliations and the topology of 3-manifolds*, J. Differential Geom. **18** (1983), no. 3, 445–503. MR **86a**:57009
- [Gab01] ———, *3 lectures on foliations and laminations on 3-manifolds*, Laminations and foliations in dynamics, geometry and topology (Stony Brook, NY, 1998), Contemp. Math., vol. 269, Amer. Math. Soc., Providence, RI, 2001, pp. 87–109. MR **2002g**:57032
- [GN95] F. J. García and A. M. Naveira, *Two remarks about foliations and minimal foliations of codimension greater than two*, Analysis and geometry in foliated manifolds (Santiago de Compostela, 1994), World Sci. Publishing, River Edge, NJ, 1995, pp. 29–38. MR **97g**:53034
- [Gro85] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), no. 2, 307–347. MR **87j**:53053
- [GS99] Robert E. Gompf and András I. Stipsicz, *4-manifolds and Kirby calculus*, American Mathematical Society, Providence, RI, 1999. MR **2000h**:57038
- [HL82] Reese Harvey and H. Blaine Lawson, Jr., *Calibrated foliations (foliations and mass-minimizing currents)*, Amer. J. Math. **104** (1982), no. 3, 607–633. MR **84h**:53095
- [KM94] P. B. Kronheimer and T. S. Mrowka, *The genus of embedded surfaces in the projective plane*, Math. Res. Lett. **1** (1994), no. 6, 797–808. MR **96a**:57073




FIGURE 10. Filling-in a logarithmic transformation

- [Kro99] P. B. Kronheimer, *Minimal genus in $S^1 \times M^3$* , Invent. Math. **135** (1999), no. 1, 45–61. MR **2000c**:57071
- [OS00] Peter Ozsváth and Zoltán Szabó, *The symplectic Thom conjecture*, Ann. of Math. (2) **151** (2000), no. 1, 93–124. MR **2001a**:57049
- [Rum79] Hansklaus Rummeler, *Quelques notions simples en géométrie riemannienne et leurs applications aux feuilletages compacts*, Comment. Math. Helv. **54** (1979), no. 2, 224–239. MR **80m**:57021
- [Sco03] Alexandru Scorpan, *Existence of foliations on 4-manifolds*, arXiv: math.GT/0302318.
- [Sul76] Dennis Sullivan, *Cycles for the dynamical study of foliated manifolds and complex manifolds*, Invent. Math. **36** (1976), 225–255. MR 55 #6440
- [Sul79] ———, *A homological characterization of foliations consisting of minimal surfaces*, Comment. Math. Helv. **54** (1979), no. 2, 218–223. MR **80m**:57022
- [Thu74] William Thurston, *The theory of foliations of codimension greater than one*, Comment. Math. Helv. **49** (1974), 214–231. MR 51 #6846

- [Thu76] W. P. Thurston, *Existence of codimension-one foliations*, Ann. of Math. (2) **104** (1976), no. 2, 249–268. MR 54 #13934
- [Thu86] William P. Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. **59** (1986), no. 339, i–vi and 99–130. MR **88h**:57014

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY,
970 EVANS HALL, BERKELEY, CA 94720

E-mail address: scorpan@math.berkeley.edu

URL: www.math.berkeley.edu/~scorpan